

## EXACT MODELS FOR ANISOTROPIC FLUID SPHERE

**K.Komathiraj**

Department of Mathematical Sciences  
Faculty of applied Sciences  
South Eastern University of Sri Lanka  
komathiraj@seu.ac.lk

### ABSTRACT

Two categories of exact solutions are found to the Einstein field equations for an anisotropic fluid sphere with a particular choice of the anisotropic factor and one of the gravitational potentials. The condition of pressure isotropy is reduced to a linear second order differential equation which can be solved in general. Consequently I can find exact solutions to the Einstein field equations corresponding to a static spherically symmetric gravitational potential in terms of elementary functions, namely polynomials and product of polynomials and algebraic functions. These solutions contain particular solutions found previously including models of isotropic relativistic spheres

**Keywords:** Exact solutions, Einstein field equations, Relativistic spheres

### INTRODUCTION

Exact solutions to the Einstein field equations with anisotropic matter have been studied by many investigators in recent years. Such Solutions for static spherically symmetric interior spacetimes are important in describing compact objects in relativistic astrophysics. Researchers have attempted to introduce different approaches of finding solutions to the field equations. Hansraj and Maharaj (2006) found solutions to the Einstein-Maxwell system with a specified form of the electrical field with isotropic pressures. These solutions satisfy a barotropic equation of state and regain the Finch and Skea (1989) model. Some of the researchers considered anisotropic pressures in the presence of the electromagnetic field with the linear equation of state of strange stars with quark matter. The approach of Esculpi and Aloma (2010) is interesting in that it utilizes the existence of a conformal symmetry in the spacetime manifold to find a solution. These exact solutions are relevant in the description of dense relativistic astrophysical objects.

In order to integrate the field equations, various restrictions have been placed by investigators on the geometry of space time and the matter content. Mainly two distinct procedures have been adopted to solve these equations for spherically symmetric static models. Firstly, the coupled differential equations are solved by computation after choosing an equation of state. Secondly, the exact Einstein solutions can be obtained by specifying the geometry and the form of the anisotropic factor. The later technique has been used by Takisa and Maharaj (2013) to produce solutions in terms of special functions and elementary functions that are suitable for the description of relativistic charged stars.

The principal objective of this work is twofold. Firstly, I seek to model a relativistic sphere with anisotropic matter which is physically acceptable. I require that the gravitational fields and matter variables are finite, continuous and well behaved in the stellar interior and the solution is stable with respect to radial perturbations. Secondly, I seek to regain an isotropic solution of Einstein field equations which satisfy the relevant physical criteria when the anisotropy factor vanishes. This ideal is not easy to achieve in practice and only a few examples with the required two features have been found thus far. The main objective of this paper is to provide systematically a solution

to Einstein equations with anisotropic matter which satisfy the above two conditions. In Section 2, the Einstein field equations for the static spherically symmetric line element with anisotropic matter is expressed as an equivalent set of differential equations utilizing a transformation. I chose particular forms for one of the gravitational potentials and the anisotropic factor, which enables me to obtain the condition of pressure anisotropy in the remaining gravitational potential. This is the master equation which determines the solvability of the entire system. It is possible to exhibit exact solutions to the Einstein field equations in terms of elementary functions. Physical properties of the solutions are briefly discussed in Section 3.

### FIELD EQUATIONS

Assume that the interior of a relativistic star should be spherically symmetric. Therefore there exists coordinates  $t$  time and  $(r, \theta, \varphi)$  spherical coordinates such that the line element is of the form

$$ds^2 = -e^{2\mu(r)} dt^2 + e^{2\lambda(r)} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \quad (1)$$

where  $\mu(r)$  and  $\lambda(r)$  are arbitrary functions related to the gravitational potentials. The system of Einstein field equations becomes for the line element (1)

$$\rho = \frac{1}{r^2}(1 - e^{-2\lambda}) + \frac{2e^{-2\lambda}}{r} \frac{d\lambda}{dr} \quad (2a)$$

$$p_r = -\frac{1}{r^2}(1 - e^{-2\lambda}) + \frac{2e^{-2\lambda}}{r} \frac{d\mu}{dr} \quad (2b)$$

$$p_t = e^{-2\lambda} \left[ \frac{d^2\mu}{dr^2} + \left(\frac{d\mu}{dr}\right)^2 + \frac{1}{r} \left(\frac{d\mu}{dr}\right) - \frac{d\mu}{dr} \frac{d\lambda}{dr} - \frac{1}{r} \left(\frac{d\lambda}{dr}\right) \right] \quad (2c)$$

$$\nabla = p_t - p_r \quad (2d)$$

The quantity  $\rho$  is the energy density,  $p_r$  is radial the pressure,  $p_t$  is the tangential pressure and  $\nabla$  is the anisotropic factor. This system governs the behaviour of the gravitational field for an anisotropic perfect fluid. To solve this system it is necessary to choose two of the variables.

In this approach  $\lambda$  and  $\nabla$  are specified. The remaining unknowns are then obtained from the rest of the system.

It is convenient at this point to introduce the choices

$$e^{2\lambda(r)} = \frac{1 - Kr^2}{1 - r^2}, \quad \nabla = \frac{\alpha Kr^2}{(1 - Kr^2)^2}$$

where  $K$  and  $\alpha$  are arbitrary constants. On using these choices, the equations (2b)-(2d) becomes

$$(1 - r^2) \left[ \frac{d^2\mu}{dr^2} + \left(\frac{d\mu}{dr}\right)^2 - \frac{1}{r} \frac{d\mu}{dr} \right] - \frac{r(1 - K)}{(1 - Kr^2)} \left( \frac{1}{r} + \frac{d\mu}{dr} \right) + (1 - K) - \frac{\alpha Kr^2}{(1 - Kr^2)^2} = 0 \quad (3)$$

which is a nonlinear differential equation.

To linearise this equation it is now convenient to introduce the transformation

$$\Psi(x) = e^{2\mu(r)}, x^2 = 1 - r^2$$

With this transformation, the nonlinear differential equation (3) becomes

$$(1 - K + Kx^2) \frac{d^2\Psi}{dx^2} - Kx \frac{d\Psi}{dx} + K(K - \alpha - 1)\Psi = 0 \quad (4)$$

Note that the Einstein system (2) implies

$$\rho = \frac{(1 - K)(3 - K + Kx^2)}{(1 - K + Kx^2)^2}$$

$$p_r = \frac{1}{(1 - K + Kx^2)} \left( -\frac{2Lx}{\Psi} \frac{d\Psi}{dx} + K - 1 \right)$$

$$p_t = \frac{1}{(1 - K + Kx^2)} \left( -\frac{2Lx}{\Psi} \frac{d\Psi}{dx} + K - 1 \right) + \frac{\alpha K(1 - x^2)}{(1 - K + Kx^2)^2}$$

In terms of the independent variable  $x$  and dependent variable  $\Psi$ . Equation (4) is the second order linear differential equation in terms of the new variables  $\Psi$  and  $x$ , and is the master equation for the system (2). Also the equation (4) to be solved to find  $\Psi$ , i.e the metric function  $\mu$ .

It is possible to express the solution of (4) in terms of special functions namely the Gegenbauer functions. However that form of the solution is not particularly useful because of the analytic complexity of the special functions involved. In addition the role of parameters of physical interest, such as the spheroidal parameter  $K$ , is lost or obscured in the representation as Gegenbauer functions. The representation of the solutions in a simple form is necessary for a detailed physical analysis. Consequently I attempt to obtain a general solution to the differential equation (4) in a series form using the method of Frobenius. Later I will indicate that it is possible to extract solutions in terms of polynomials and algebraic functions for particular parameter values as demonstrated by Komathiraj and Maharaj (2010).

As the point  $x = 0$  is the regular point of the differential equation (4), there are two linearly independent solutions. Thus the general solution to (4) can be assumed by the method of Frobenius as

$$\Psi(x) = \sum_{i=0}^{\infty} a_i x^i$$

where  $a_i$  are constants to be determined.

Two classes of solutions in terms of elementary functions for (4) are possible from the above series form. The first category of solutions for  $\Psi(x)$  is given by

$$\Psi_1(x) = A \sum_{j=0}^n (-\beta)^j \frac{(n+j-2)!}{(n-j)!(2j)!} x^{2j} \\ + B(1-K+Kx^2)^{\frac{3}{2}} \sum_{j=0}^{n-2} (-\beta)^j \frac{(n+j)!}{(n+j-2)!(2j+1)!} x^{2j+1}$$

for the values

$$\beta = 4 - \frac{4}{4n(n-1) - \alpha} \\ K - \alpha = [2 - (2n-1)^2]$$

The second category of solution for  $\Psi(x)$  has the form

$$\Psi_2(x) = A \sum_{j=0}^n (-\gamma)^j \frac{(n+j-1)!}{(n-j)!(2j+1)!} x^{2j+1} \\ + B(1-K+Kx^2)^{\frac{3}{2}} \sum_{j=0}^{n-1} (-\gamma)^j \frac{(n+j)!}{(n-j-1)!(2j)!} x^{2j}$$

with the values

$$\gamma = 4 - \frac{4}{4n^2 - 1 - \alpha} \\ K - \alpha = 2(1 - 2n)^2$$

In the above two categories of solutions  $A$  and  $B$  are arbitrary constants.

## DISCUSSION

One of the original reasons for studying anisotropic matter was to generate models that permit redshifts higher than the critical redshift of isotropic matter. Observational results indicate that certain isolated objects have redshifts higher than critical redshift of isotropic matter. I have found new solutions to the Einstein field equations for an anisotropic fluid sphere by utilizing the method of Frobenius for an infinite series; a particular form for one of the gravitational potentials was assumed and the anisotropic factor was specified. These solutions are given in terms of special functions. For particular values of the parameters involved it is possible to write the solution in terms of elementary functions: polynomials and products of polynomials and algebraic functions. The anisotropic factor may vanish in the solutions anisotropic solutions can be regained. Thus this approach has the advantage of necessarily containing anisotropic stellar solution.. The simple form of the solutions found facilitates the analysis of the physical features of an anisotropic fluid sphere. The gravitational potentials are finite at the centre  $r = 0$  and at the boundary  $r = R$ . These functions are continuous and well behaved in the interior of the relativistic star. The radial pressure is continuous and well behaved in the interior of the star. Also the radial pressure is greater than zero in the interval  $(0,R)$ , regular at the centre, and vanishes at the boundary. In general the tangential pressure is not zero at the boundary of the star which is different from the radial pressure.

## CONCLUSION

The main objective of this work was to find exact solutions to the anisotropic fluid sphere which can be used to describe a relativistic dense star. Solutions of the complicated system of nonlinear partial differential equations were sought by

specifying physically reasonable forms for one of the gravitational potentials and the anisotropic factor. A number simple solutions to the system, which we believe to be physically reasonable, were obtained explicitly in terms of elementary functions. It was also possible to find other categories of solutions by specifying other types of spatial geometries.

#### REFERENCES

- Hansraj S and S. D. Maharaj S. D. (2006), Charged Analogue of Finch-Skea Stars, *Int. J. Mod. Phys. D* 15. p. 1311-1327
- Finch M. R and Skea J. E. F. (1989), A realistic stellar model based on an ansatz of Duorah and Ray, *Class. Quantum. Grav.* 6. p. 467- 476
- M. Esculpi and Aloma E. (2010), Conformal anisotropic relativistic charged fluid spheres with the linear equation of state, *Eur. Phys. J. C* 67. p. 521- 532
- Mafa Takisa P and Maharaj S. D. (2013), Compact models with regular charge distributions, *Astrophys. Space Sci.* 343. p. 569-577
- Komathiraj K and Maharaj S. D. (2010), A class of charged relativistic spheres, *Math.Com. App.* 15. p. 665-673